

Corrigenda: Daniel CA, Midgley JM, Villet MH (2017) Determination of species and instars of the larvae of the Afrotropical species of *Thanatophilus* Leach, 1817 (Coleoptera, Silphidae). African Invertebrates 58(2): 1-10. <https://doi.org/10.3897/afrinvertebr.58.12966>

C.A. Daniel¹, J.M. Midgley^{1,2,3}, M.H. Villet¹

1 Southern African Forensic Entomology Research Laboratory, Department of Zoology and Entomology, Rhodes University, P.O. Box 94, Grahamstown, 6140 South Africa **2** Department of Entomology and Arachnology, Albany Museum, Somerset Street, Grahamstown, 6139 South Africa **3** Current address: KwaZulu-Natal Museum, Private Bag 9070, Pietermaritzburg, 3200, South Africa

Corresponding author: J.M. Midgley (johnmidge@gmail.com)

Academic editor: Y. Mutafchiev | Received 1 September 2017 | Accepted 1 September 2017 | Published 25 October 2017

<http://zoobank.org/C38390BA-1245-4C4D-9F5C-3F63C5E03197>

Citation: Daniel CA, Midgley JM, Villet MH (2017) Corrigenda: Daniel CA, Midgley JM, Villet MH (2017) Determination of species and instars of the larvae of the Afrotropical species of *Thanatophilus* Leach, 1817 (Coleoptera, Silphidae). African Invertebrates 58(2): 1-10. <https://doi.org/10.3897/afrinvertebr.58.12966>. African Invertebrates 58(2): 53–53. <https://doi.org/10.3897/AfrInvertebr.58.20751>

After the publication of this article, it was brought to our attention that there was an error in the caption for Figure 2. The caption should read:

Figure 2. Right mandible in dorsal view and mesothoracic spiracle of mature Afrotropical *Thanatophilus* larvae. The arrows indicate species-specific diagnostic characteristics: *T. micans* has two ventrolateral mandibular setae and two spiracular setae, while *T. mutilatus* has three ventrolateral mandibular setae and one spiracular seta.